

Temperature transfer functions based on freshwater testate amoebae from China

Jean Claude Ndayishimiye^{a,b,1}, Lihua Ju^{a,b,1}, Hongkai Li^c, Xiangdong Yang^d, Zhengwen Liu^d, Jun Yang^{a,*}

^a*Aquatic EcoHealth Group, Key Laboratory of Urban Environment and Health, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, PR China*

^b*University of Chinese Academy of Sciences, Beijing 100049, PR China*

^c*State Environmental Protection Key Laboratory of Wetland Ecology and Vegetation Restoration, Northeast Normal University, Changchun 130024, PR China*

^d*State Key Laboratory of Lake Science and Environment, Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences, Nanjing 210008, PR China*

Abstract

Globally, lakes and reservoirs ecosystems are integral parts of ecological processes. Nevertheless, global warming is rapidly changing their function and sustainability, especially in the populated area of the northern hemisphere. Here we examined testate amoebae community and 10 environmental variables from 51 lakes and reservoirs across China and developed testate amoebae transfer functions for temperature based on both abundance- and biomass-datasets. A total of 169 testate amoebae taxa were identified. Our partial CCA analyses revealed that water temperature explained 5.15% ($P = 0.006$) and 5.57% ($P = 0.008$) of the total variance in testate amoebae abundance and biomass, respectively. The WA-PLS was the best model in abundance-based temperature transfer function (RMSEP = 2.87 °C, $R^2 = 0.60$), whereas the MAT proved to be the best model for biomass-based temperature transfer function (RMSEP = 3.34 °C, $R^2 = 0.67$). The application of all models should be carried out with suitable precautions. Our results suggested that freshwater testate amoebae could contribute to a better understanding of the ecological integrity and its vulnerability in inland aquatic environments.

*Corresponding author

European Journal of Protistology

Received 15 March 2018/Accepted 22 March 2019

<https://doi.org/10.1016/j.ejop.2019.03.003>

0932-4739/© 2019 Elsevier GmbH. All rights reserved.